

Version: 2025

TDS2

## **BROMUS BLOCCO 10kg**

"Rodent control"



# **BROMUS BLOCCO 10kg**

Codice prodotto: 2450631A

Category: Biocide

Ready-to-use rodenticide bait in paraffin block form for trained professionals.

## **Italian Ministry of Health Registration:**

IT/2014/00198/AUT

#### Formula:

Block

### **Composition:**

#### 100g of product contains:

Brodifacoum: 0.05g

Denatonium Benzoate: 0.01g

### **Description:**

BROMUS BLOCK is a ready-to-use rodenticide bait based on an anticoagulant active substance, called Brodifacoum, effective against the house mouse (*Mus musculus*), brown or common rat (*Rattus norvegicus*) and black rat (*Rattus rattus*), even after a single ingestion. The formulation consists of a paraffin block that is palatable for the indicated rodent species. The product does not alert or arouse suspicion in other rodent populations. BROMUS BLOCK contains a bitter substance to prevent accidental ingestion by children. BROMUS BLOCK can be used in and around industrial rural, civil and residential buildings (including warehouses, stores, ship holds), cellars, garages, storage rooms, gardens and outdoor areas. **TARGET ORGANISMS** Young and adult house mice (*Mus musculus*), young and adult brown rats (*Rattus norvegicus*), young and adult black rats (*Rattus rattus*). **METHOD OF USE** BROMUS BLOCK is a ready-to-use bait in the form of a paraffin block, to be used in tamper-proof bait containers or in covered bait points protected from weathering, ingestion by non-target species and dispersal in the environment. Aroma: berries

#### Method of use:

Use inside buildings – For control of house mouse (*Mus musculus*), brown rat (*Rattus norvegicus*) and black rat (*Rattus rattus*). In case of high infestation: place 100 g of BROMUS

BLOCK per container, amounting to 5 baits in ready-to-use paraffin blocks of 20 grams each. In the case of low infestation:



Version: 2025

TDS2

## **BROMUS BLOCCO 10kg**

"Rodent control"


place 60 g of BROMUS BLOCK per container, equal to 3 baits in ready-to-use paraffin blocks of 20 g each. House mouse ( *Mus musculus*) – Place 40 g of BROMUS BLOCK per container, equivalent to 2 baits in ready-to-use paraffin blocks of 20 g each. Remove residual product at the end of the treatment.

Follow any further instructions required by relevant good practice. Where possible, before treatment, inform possible bystanders (those frequenting the treated area and its surroundings) of the ongoing pest control campaign. Consider preventive control measures to encourage uptake and reduce the likelihood of re-infestation (e.g. blocking possible openings, removing potential food and water sources as much as possible). In order to reduce the risk of secondary poisoning, search for and collect dead rodents during treatment and at frequent intervals, in accordance with relevant good practice recommendations. Do not use the product as a permanent bait for preventing rodent infestations or monitoring rodent activity. Do not use the product in treatments where 'pulsed baiting' is used.

When placing bait points near water drainage systems, prevent the bait from coming into contact with water. **Use outdoors, around buildings** – For the control of House mice (*Mus musculus*), Brown rats (*Rattus norvegicus*) and Black rats (*Rattus rattus*).

Brown rat (*Rattus norvegicus*) and Black rat (*Rattus rattus*) – In the case of high infestation: place 100 g of BROMUS BLOCK per container, equal to 5 baits in paraffin blocks of 20 g each. In the case of low infestation: place 60 g of BROMUS BLOCK per container, equal to 3 baits in ready-to-use paraffin blocks of 20 g each. House mouse (*Mus musculus*) – Place 40 g of BROMUS BLOCK per container, equal to 2 baits in ready-to-use paraffin blocks of 20 grams each. Remove residual product at the end of the treatment. For applications in covered and protected bait points: for outdoor use, bait points should be covered and placed in strategic locations to minimise exposure to non-target species.

Do not use this product directly in rodent nests. Where possible, before treatment, inform possible bystanders (those frequenting the treated area and its surroundings) of the ongoing pest control campaign. Consider preventive control measures to encourage uptake of the product and reduce the likelihood of re-infestation (e.g. blocking possible openings, removing potential food and water sources as much as possible). In order to reduce the risk of secondary poisoning, search for and collect dead rodents during treatment and at frequent intervals, in accordance with relevant good practice recommendations. When placing bait points near surface water (e.g. rivers, ponds, waterways, dams, irrigation canals) or water drainage systems, prevent the bait from coming into contact with water. Do not use the product as a permanent bait for preventing rodent infestations or monitoring rodent activity. Do not use the product in treatments where 'pulsed baiting' is



Version: 2025

TDS2

## **BROMUS BLOCCO 10kg**

"Rodent control"

## Warnings:

Before use, read and follow the instructions on the product packaging as well as the information accompanying the product or provided at the point of sale.

Before positioning baits, carry out a survey of the infested area and an on-site assessment in order to identify the rodent species, places of activity and determine the probable cause and extent of the infestation.

Remove food sources easily accessible to rodents (e.g. spilled grain or food scraps). Also, do not clean up the infested area just before treatment, as this only disturbs the rodent population and makes it more difficult to gain acceptance of the bait. The product should only be used within the framework of an integrated pest management (IPM) system, which includes, among other things, hygiene measures and, where possible, physical control methods.

Consider preventive control measures (e.g. closing holes, removal, as far as possible, of all food and water sources) in order to control the amount of product consumed and reduce the possibility of re-infestation.

Containers with bait must be placed in the immediate vicinity of places where rodent activity has been previously verified (e.g. walkways, nests, livestock entrance fences, openings, lairs, etc.).

#### Unit of sales:

10kg box

### **Disposal information:**

Ricicliamo insieme			
Element	Material		Further information
Boxes Ties	Cardboard Composite material	PAP20 C/PVC 92	Paper recycling Undiferentiated waste collection
Envelope	Plastic	LDPE 4	he packaging contains a rodenticide; please check the regulations of your municipality before disposal. Recycling: check the regulations of your municipality

FOLLOW THE INSTRUCTIONS OF YOUR MUNICIPALITY