



## PIRETRO SAFE EC® 1I

**Codice prodotto: 0860010**

**Category: PMC**

Concentrated emulsion Pyrethrins based insecticide

**Italian Ministry of Health Registration:**

n. 15288

**Formula:**

Emulsifiable concentrate

**Composition:**

**100g of product contains:**

Chrysanthemum cinerariaefolium extract: 1.5g  
(Equivalent to Extract of Pyrethrum 50%): 3.0g  
Piperonyl Butoxide: 12.0g  
Coformulants: 30.1g

**Description:**

PIRETRO SAFE EC kills insects by contact and ingestion. The insect mortality intervenes in 24 hours after contact with the product, for a minimum duration of 15 days. PIRETRO SAFE EC is effective against flying and crawling insects, it may especially be used:

- in livestock buildings to kill houseflies (*Musca domestica*)
- public parks, perimeter zones of houses and factories to kill mosquitoes (*Aedes spp.* and *Culex spp.*)
- in buildings, plants and food industries especially to kill ants (*Lasius spp.*), indian mealmoth (*Plodia interpunctella*), mediterranean flour moth (*Ephestia kuehniella*), the flour beetle (*Tribolium spp.*), the granary weevil (*Sitophilus spp.*), cockroaches (*Blattella germanica*, *Blatta orientalis* and *Periplaneta americana*), lesser grain beetle (*Rhyzopertha Dominica*) and sawtoothed grain beetle (*Oryzaephilus surinamensis*).

**Method of use:**

PIRETRO SAFE EC must be used at a concentration of 4 to 5% (dilute 400 to 500ml of product in 10 liters of water). Use a liter of solution to treat 10 to 20 m<sup>2</sup> of walls / surfaces treated or 400 m<sup>3</sup>.

Use diluted solution immediately. The solution should be sprayed with a handsprayer or a cold (Nebulo) or hot nebulizer. When applied by nebulization in closed rooms, a breakdown of two hours is recommended.

For use against the flour beetle (*Tribolium* spp.) The application must be made by a cold spray. For use in livestock buildings, it is recommended to apply the product on vertical surfaces. When applied by spraying, respect a period of re-entry of animals in treated areas at least an hour.

**User:**

Professional

**Warnings:**

Do not contaminate foodstuffs, beverages or containers for them when using the spray. When used in food processing plants, the product must not come into direct or indirect contact with food. In case of use in animal shelters, keep animals away during treatment.

**Unit of sales:**

Bottle of 1 liter

**Notes:**

Medical Device with insecticidal action, use with caution. Always read the label and product information before use.

Attention is drawn to the hazard warnings and symbols on the label.

The text in the specifications corresponds to the label approved by the Ministry of Health.