



## BROMUS BLOCK for trained professionals

**Codice prodotto: 2450631**

**Category: Biocide**

Ready-to-use rodenticide bait in the form of a paraffin wax block for professionals.

**Italian Ministry of Health Registration:**

IT/2014/00198/AUT

### Description:

BROMUS BLOCK is a ready-to-use rodenticidal bait based on an anticoagulant active substance, called Brodifacoum, effective against house mice (*Mus musculus*), common rats (*Rattus norvegicus*) and black rats (*Rattus rattus*), even after a single ingestion. The formulation consists of a paraffin wax block that is palatable to the indicated rodent species. The product does not alert or arouse suspicion in other rodent populations.

BROMUS BLOCK contains a bitter substance to prevent accidental ingestion by children.

BROMUS BLOCK can be used in and around industrial buildings (including warehouses, stores, ship holds), rural, civil, residential, cellars, garages, storage rooms, gardens and outdoor areas.

### Method of use:

BROMUS BLOCK is a ready-to-use bait in the form of a paraffin wax block, to be used exclusively in tamper-proof, properly marked bait containers, protected from the weather, ingestion by non-target species and dispersal in the environment.

For use indoors: for the control of house mice (*Mus musculus*), common rats (*Rattus norvegicus*) and black rats (*Rattus rattus*):

Common rats (*Rattus norvegicus*) and black rats (*Rattus rattus*):

in case of high infestation, place 100 gr of BROMUS BLOCK per container, corresponding to 5 baits in ready-to-use paraffin wax blocks of 20 grams each. in case of low infestation, place 60 gr of BROMUS BLOCK per container, corresponding to 3 baits in ready-to-use paraffin wax blocks of 20 grams each.

House mice (*Mus musculus*): place 40 g of BROMUS BLOCK per container, corresponding to 2 baits in ready-to-use paraffin wax blocks of 20 g each.

Remove any remaining product at the end of the treatment.

Follow any further instructions as required by relevant good practice.

Whenever possible, before treatment, inform possible bystanders (e.g. those frequenting the treated area and its surroundings) of the ongoing pest control campaign. Consider preventive control measures to encourage uptake and reduce

the likelihood of re-infestation (e.g. blocking possible openings, removing potential food and water sources as much as possible).

In order to reduce the risk of secondary poisoning, search for and collect dead rodents during treatment and at frequent intervals, in accordance with relevant recommendations of good practice.

Do not use the product as a permanent bait for the prevention of rodent infestations or for monitoring rodent activity.

Do not use the product in treatments where 'pulsed baiting' is used.

When placing bait points near water drainage systems, prevent the bait from coming into contact with water.

Outdoor use, around buildings. For the control of house mice (*Mus musculus*), common rats (*Rattus norvegicus*) and black rats (*Rattus rattus*).

Common rats (*Rattus norvegicus*) and black rats (*Rattus rattus*):

Place 100 gr of BROMUS BLOCK per container, corresponding to 5 baits in ready-to-use paraffin wax blocks of 20 grams each.

in case of low infestation: place 60 gr of BROMUS BLOCK per container, corresponding to 3 baits in ready-to-use paraffin wax blocks of 20 grams each.

House mice (*Mus musculus*): place 40 gr of BROMUS BLOCK per container, corresponding to 2 baits in ready-to-use paraffin wax blocks of 20 g each.

Remove residual product at the end of the treatment.

For covered and protected bait point applications: for outdoor use, the bait points should be covered and placed in strategic locations to minimise exposure to non-target species.

Do not use this product directly in nests.

Whenever possible, before treatment, inform potential bystanders (e.g. those frequenting the treated area and its surroundings) of the ongoing pest control campaign. Consider preventive control measures to encourage uptake and reduce the likelihood of re-infestation (e.g. blocking possible openings, removing potential food and water sources as much as possible).

In order to reduce the risk of secondary poisoning, search for and collect dead rodents during treatment and at frequent intervals, in accordance with relevant recommendations of good practice.

If bait points are placed near surface water (e.g. rivers, ponds, water channels, dams, irrigation canals) or water drainage systems, prevent the bait from coming into contact with the water.

Do not use the product as a permanent bait for the prevention of rodent infestations or for monitoring rodent activity. Do not use the product in treatments where 'pulsed baiting' technique is used.



Version: 2024

TDS2

**BROMUS BLOCK for trained professionals**

**"Rodent control"**

**Rodenticides**

**User:**

Professional

**Warnings:**

Before use, read and follow the instructions on the product packaging as well as the information accompanying the product or provided at the point of sale.

Prior to positioning the bait, check the infested area and carry out an on-site assessment to identify rodent species, locations of activity and determine the probable cause and extent of infestation.

Remove food sources within easy reach of rodents (e.g. spilled grain or food waste).

Moreover, do not clean up the infested area just prior to treatment, as this will only disturb the rodent population and make it more difficult to gain acceptance of the bait.

The product must only be used as part of an integrated pest management (IPM) system, which includes, inter alia, hygiene measures and, where possible, physical control methods.

Consider preventive control measures to encourage uptake and reduce the likelihood of re-infestation (e.g. blocking possible openings, removing potential food and water sources as much as possible).

Bait containers must be placed in the immediate vicinity of places where rodent activity has been previously verified (e.g. walkways, nests, livestock entrance fences, openings, burrows, etc.).

**Unit of sales:**

10 kg box

**Notes:**

Use biocides with caution. Always read the label and product information before use.

Attention is drawn to the hazard warnings and symbols on the label.

The text in the specifications corresponds to the label approved by the Ministry of Health.